

## **Original references of Basalts used to calculate Tp in this study**

### **Greenland Craton**

T. Komiya, S. Maruyama, T. Hirata, H. Yurimoto, S. Nohda, Geochemistry of the oldest MORB and OIB in the Isua Supracrustal Belt, southern West Greenland: implications for the composition and temperature of early Archean upper mantle, *Island Arc* 13 (2004) 47–72.

J.C. Ordóñez-Calderón, A. Polat, B.J. Fryer, P.W.U. Appel, J.A.M. van Gool, Y. Dilek, J.E. Gagnon, Geochemistry and geodynamic origin of the Mesoarchean Ujarassuit and Ivisaartoq greenstone belts, SW Greenland, *Lithos* 113 (2009) 133–157.

A. Polat, R. Frei, P.W.U. Appel, Y. Dilek, B. Fryer, J.C. Ordóñez-Calderón, Z. Yang, The origin and compositions of Mesoarchean oceanic crust: evidence from the 3075 Ma Ivisaartoq greenstone belt, SW Greenland, *Lithos* 100 (2008) 293–321.

A. Polat, P.W.U. Appel, R. Frei, Y. Pan, Y. Dilek, J.C. Ordóñez-Calderón, B. Fryer, J.A. Hollis, J.G. Raith, Field and geochemical characteristics of the Mesoarchean (~ 3075 Ma) Ivisaartoq greenstone belt, southern West Greenland: Evidence for seafloor hydrothermal alteration in supra-subduction oceanic crust, *Gondwana Research* 11 (2007) 69–91.

H. Furnes, M. de Wit, H. Staudigel, M. Rosing, K. Muehlenbachs, A vestige of Earth's oldest ophiolite, *Science* (1979) 315 (2007) 1704–1707.

A. Polat, A.W. Hofmann, C. Münker, M. Regelous, P.W.U. Appel, Contrasting geochemical patterns in the 3.7–3.8 Ga pillow basalt cores and rims, Isua greenstone belt, Southwest Greenland: implications for postmagmatic alteration processes, *Geochim Cosmochim Acta* 67 (2003) 441–457.

### **Superior and Wyoming craton, USA and Canada**

J. Dostal, W.U. Mueller, Deciphering an Archean mantle plume: Abitibi greenstone belt, Canada, *Gondwana Research* 23 (2013) 493–505.

A. Polat, R. Kerrich, Archean greenstone belt magmatism and the continental growth–mantle evolution connection: constraints from Th–U–Nb–LREE systematics of the 2.7 Ga Wawa subprovince, Superior Province, Canada, *Earth Planet Sci Lett* 175 (2000) 41–54.

P. Hollings, R. Kerrich, Geochemical systematics of tholeiites from the 2.86 Ga Pickle Crow assemblage, northwestern Ontario: arc basalts with positive and negative Nb–Hf anomalies, *Precambrian Res* 134 (2004) 1–20.

A. Polat, R. Kerrich, D.A. Wyman, The late Archean Schreiber–Hemlo and White River–Dayohessarah greenstone belts, Superior Province: collages of oceanic plateaus, oceanic arcs, and subduction–accretion complexes, *Tectonophysics* 289 (1998) 295–326.

S.B. Goldstein, D. Francis, The petrogenesis and mantle source of Archean ferropicrites from the Western Superior Province, Ontario, Canada, *Journal of Petrology* 49 (2008) 1729–1753.

J. Fan, R. Kerrich, Geochemical characteristics of aluminum depleted and undepleted komatiites and HREE-enriched low-Ti tholeiites, western Abitibi greenstone belt: A heterogeneous mantle plume-convergent margin environment, *Geochim Cosmochim Acta* 61 (1997) 4723–4744.

S.-J. Barnes, The petrography and geochemistry of komatiite flows from the Abitibi Greenstone Belt and a model for their formation, *Lithos* 18 (1985) 241–270.

### **Barberton craton, South Africa**

H. Furnes, B. Robins, M.J. de Wit, Geochemistry and petrology of lavas in the upper Onverwacht suite, Barberton Mountain Land, South Africa, *South African Journal of Geology* 115 (2012) 171–210.

### **Pilbara craton, Australia**

Y. Kato, K. Nakamura, Origin and global tectonic significance of Early Archean cherts from the Marble Bar greenstone belt, Pilbara Craton, Western Australia, *Precambrian Res* 125 (2003) 191–243.

M.G. Green, P.J. Sylvester, R. Buick, Growth and recycling of early Archaean continental crust: geochemical evidence from the Coonterunah and Warrawoona Groups, Pilbara Craton, Australia, *Tectonophysics* 322 (2000) 69–88.

### **Yilgarn craton, Australia**

N. Said, R. Kerrich, D. Groves, Geochemical systematics of basalts of the Lower Basalt Unit, 2.7 Ga Kambalda Sequence, Yilgarn craton, Australia: plume impingement at a rifted craton margin, *Lithos* 115 (2010) 82–100.

N. Said, R. Kerrich, Geochemistry of coexisting depleted and enriched Paringa Basalts, in the 2.7 Ga Kalgoorlie Terrane, Yilgarn Craton, Western Australia: Evidence for a heterogeneous mantle plume event, *Precambrian Res* 174 (2009) 287–309.

### **Baltic craton, Karelian Province, Finland**

P. Hölttä, E. Heilimo, H. Huhma, A. Kontinen, S. Mertanen, P. Mikkola, J. Paavola, P. Peltonen, J. Semprich, A. Slabunov, The Archaean of the Karelia province in Finland, (2012).

A.A. Shchipansky, A. v Samsonov, E. v Bibikova, I.I. Babarina, A.N. Konilov, K.A. Krylov, A.I. Slabunov, M.M. Bogina, 2.8 Ga boninite-hosting partial suprasubduction zone ophiolite sequences from the North Karelian greenstone belt, NE Baltic Shield, Russia, *Developments in Precambrian Geology* 13 (2004) 425–486.

H. Papunen, T. Halkoaho, E. Luukkonen, Archaean evolution of the tipasjärvi-Kuhmo-Suomussalmi greenstone complex, Finland, *Bulletin of the Geological Survey of Finland* (2009) 1–68.

### **Dharwar craton, India**

N.M. Noack, R. Kleinschrodt, M. Kirchenbaur, R.O.C. Fonseca, C. Müunker, Lu–Hf isotope evidence for Paleoproterozoic metamorphism and deformation of Archean oceanic crust along the Dharwar Craton margin, southern India, *Precambrian Res* 233 (2013) 206–222.

S.M. Naqvi, R.M.K. Khan, C. Manikyamba, M.R. Mohan, T.C. Khanna, Geochemistry of the NeoArchaean high-Mg basalts, boninites and adakites from the Kushtagi-Hungund greenstone belt of the

Eastern Dharwar Craton (EDC); implications for the tectonic setting, *J Asian Earth Sci* 27 (2006) 25–44.  
<https://doi.org/10.1016/j.jseas.2005.01.006>.

## Proterozoic

### Flin Flon and Cape Smith, Canada

M.G. Babechuk, B.S. Kamber, An estimate of 1.9 Ga mantle depletion using the high-field-strength elements and Nd–Pb isotopes of ocean floor basalts, *Flin Flon Belt, Canada, Precambrian Res* 189 (2011) 114–139.

A. Hynes, D.M. Francis, A transect of the early Proterozoic Cape Smith foldbelt, New Quebec, *Tectonophysics* 88 (1982) 23–59.

C. Picard, D. Giovenazzo, D. Lamothe, Geotectonic evolution by asymmetric rifting of the Proterozoic Cape Smith Belt, New Quebec, *Geoscience Canada* 16 (1989) 130–134.

### Phulad Ophiolite, India

A.M. Volpe, J.D. Macdougall, Geochemistry and isotopic characteristics of mafic (Phulad Ophiolite) and related rocks in the Delhi Supergroup, Rajasthan, India: implications for rifting in the Proterozoic, *Precambrian Res* 48 (1990) 167–191.